

Black hole thermodynamics with generalized uncertainty principle

Li Xiang* and X. Q. Wen

Center for Relativistic Astrophysics and High Energy Physics,

Department of Physics, Nanchang University,

Nanchang, 330031, Jiangxi province, P. R. China

Abstract

In the standard viewpoint, the temperature of a stationary black hole is proportional to its surface gravity, $T_H = \hbar\kappa/2\pi$. This is a semiclassical result and the quantum gravity effects are not taken into consideration. This Letter explores a unified expression for the black hole temperature in the sense of a generalized uncertainty principle(GUP). Our discussion involves a heuristic analysis of a particle which is absorbed by the black hole. Besides a class of static and spherically symmetric black holes, an axially symmetric Kerr-Newman black hole is considered. Different from the existing literature, we suggest that the black hole's irreducible mass represent the characteristic size in the absorption process. The information capacity of a remnant is also discussed by Bousso's D-bound in de Sitter spacetime.

Keywords: generalized uncertainty principle, black hole temperature, irreducible mass, remnant.

PACS numbers: 04.70.Dy, 04.70.-s

*xiang.lee@163.com

1 Introduction

Heisenberg's uncertainty relation is one of the fundamental principles of quantum mechanics. This principle only meets the quantum effects of matters, and it does not directly describe the quantum fluctuations of spacetimes. However, many efforts have shown that Heisenberg's principle may suffer a modification[1]-[10], in the context of quantum gravity. Concretely, a generalized uncertainty principle(GUP) reads

$$\Delta x \geq \frac{\hbar}{\Delta p} + \frac{\alpha}{\hbar} \Delta p, \quad (1)$$

where $\alpha \sim G$. The second term on the r.h.s means a new duality, which is firstly related to the spacetime uncertainty principle[1, 3] and the scattering amplitude of high energy string[2]. This term is also attributed to gravity in some gedanken experiments[4, 5, 7, 9]. Different from Heisenberg's principle, GUP restricts the shortest distance that we can probe(i.e. $\Delta x \geq 2\sqrt{\alpha} \sim l_p$). This agrees with the belief that Planck length is a fundamental scale in quantum gravity.

Since uncertainty principle is of great importance to quantum physics, GUP has caused extensive interests and arguments. In particular, GUP's effects on the thermodynamics of a Schwarzschild black hole have been discussed by a heuristic method[10]. The crucial idea therein is that Δx and Δp are identified as the black hole's size and temperature respectively. An interesting result is that the black hole mass is not allowed to be less than a scale of order Planck mass, which suggests a black hole remnant. Although GUP's impacts on black hole thermodynamics have been discussed in the literature[10]-[25], a universal expression is still absent.

In the semiclassical framework, Hawking temperature of a stationary black hole is proportional to the surface gravity, i.e.

$$T_H = \frac{\hbar \kappa}{2\pi}, \quad (2)$$

where Planck constant reveals the quantum nature of black hole radiation. In the Bekenstein's original work[26], Heisenberg's uncertainty principle is crucial to the linear relation between Hawking temperature and surface gravity.¹ In our opinion, GUP changes the semiclassical framework to a certain context, and the semiclassical black hole temperature (2) should suffer a modification. How the expression (2) is corrected by GUP? As an extension of Ref.[10], this research explores a revised temperature expression which is expected to be valid for more

¹This point is also stressed in Ref.[28], where the linear relation $T_H \sim \hbar \kappa$ can be obtained by another heuristic method via Heisenberg's uncertainty principle.

general black holes. We discuss a class of static and spherically symmetric black holes, as well as a Kerr-Newman black hole. The temperatures of these black holes have the same form. The information capacity of a black hole remnant is discussed in de Sitter spacetime, in terms of a Bousso's D-bound. We follow the Bekenstein's original work[26], and analyze a gedanken experiment that a neutral particle just outside the horizon is absorbed by the black hole. This Letter takes the units $G = c = k_B = 1$.

2 Black holes thermodynamics: a heuristic analysis

2.1 Brief review

This subsection gives a brief review of the basis for the further discussion. Let us start with the first law of black hole mechanics[26, 29]

$$dM = \frac{\kappa}{8\pi}dA + \sum_i Y_i dy_i, \quad (3)$$

where the terms $\sum_i Y_i dy_i$ represent the work done on the black hole by an external agent. y_i are the black hole's variables such as electronic charge and angular momentum; Y_i are the generalized forces corresponding to the variables y_i , e.g. electrostatic potential and angular velocity. The above formula is a result of classical general relativity. However, it has been endowed with thermodynamic meaning since Hawking radiation was discovered, i.e.

$$dM = TdS + \sum_i Y_i dy_i. \quad (4)$$

Corresponding to the standard temperature (2), the black hole entropy is expressed as $S_{BH} = (4\hbar)^{-1}A$, i.e. the so-called Bekenstein-Hawking entropy. However, this simple relation is a semiclassical result. In more general situations, the entropy of a black hole is assumed to be a function of its area[26], $S = S(A)$. Following from (3) and the definition of thermodynamics, the temperature is expressed as

$$T = \left(\frac{\partial M}{\partial S} \right)_{y_i} = \frac{dA}{dS} \times \left(\frac{\partial M}{\partial A} \right)_{y_i} = \frac{dA}{dS} \times \frac{\kappa}{8\pi}, \quad (5)$$

where the variables y_i are fixed. The temperature expression is determined by the relation between the entropy and area. In order to find the concrete form of $S(A)$, we consider a particle captured by the black hole. When the particle disappears, on one hand, its information is lost to an observer outside the horizon; on the other hand, the smallest increase in the area of a Kerr-Newmann black hole is given by[26]

$$\Delta A \sim b\mu, \quad (6)$$

where b and μ are the particle's size and mass, respectively. Identifying the loss of information with the increase of black hole entropy, we obtain

$$\Delta S \simeq \frac{dS}{dA} \Delta A.$$

According to information theory, the loss of information is one bit at least, i.e. $(\Delta S)_{min} = \ln 2$. The next step is to work out the differential relation dS/dA via (6). For a classical particle(point-like object), $(\Delta A)_{min} = 0$. However, in quantum mechanics, a particle is described by a wave packet and a definite trajectory does not exist. The width of wave packet is defined as the standard deviation of x distribution(i.e. the position uncertainty), which can be interpreted as the characteristic size of the particle($b \sim \Delta x$). Furthermore, the momentum uncertainty is not allowed to be greater than the mass ($\Delta p \leq \mu$), in the process of measuring the particle's position. Otherwise the relativistic effects lead to the creation of a partner of the particle and make the measurement meaningless. Thus the expression (6) is deduced to

$$\Delta A \sim b\mu \geq \Delta x \Delta p. \quad (7)$$

The smallest increase in area cannot be arbitrarily small and it is restricted by the uncertainty relation of quantum mechanics. In the Bekenstein's insightful work, Heisenberg principle is utilized to identify the particle's size with the Compton wavelength of itself, and then the minimum increase in horizon area is given by $\Delta A \sim l_p^2$. This results in

$$\frac{\Delta S}{\Delta A} = const,$$

which means the linear relation between the black hole entropy and the horizon area. GUP will correct the Bekenstein's result. Substituting (1) into (7), we have

$$\Delta A \geq \gamma_1 \hbar \left[1 + \frac{\alpha}{\hbar^2} (\Delta p)^2 \right], \quad (8)$$

where γ_1 is a calibration factor. The minimum increase in area, $(\Delta A)_{min}$, is determined by the smallest uncertainty of momentum. Following from (8), $(\Delta A)_{min}$ would be a constant if $\Delta p \rightarrow 0$. At a first glimpse, there seems to be no correction to the Bekenstein's result. However, $\Delta p \rightarrow 0$ means $\Delta x \rightarrow \infty$. For a particle captured by black hole, Δp is not allowed to be arbitrarily small, since the particle is confined within a finite region and Δx is finite. $(\Delta A)_{min}$ is therefore no longer a constant, which results in some corrections to the linear relation between entropy and area. In the following subsections, a static and spherically symmetric black hole as well as an axially symmetric Kerr-Newman black hole are discussed respectively.

2.2 A class of static and spherically black holes

We consider a static and spherical black hole as follows

$$ds^2 = -F(r)dt^2 + F^{-1}(r)dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2),$$

where the horizon is located by $F(r_0) = 0$. The above line element describes a class of static and spherically symmetric black holes, such as Schwarzschild, Reissner-Nordström and their partners in (anti-)de Sitter spacetime. When a particle is captured by black hole, the position uncertainty should not be greater than a specific scale. This characteristic size, for a static and spherically symmetric black hole, is identified with the twice radius of horizon,² i.e.

$$2r_0 \geq \Delta x \geq \frac{\hbar}{\Delta p} + \frac{\alpha}{\hbar} \Delta p, \quad (9)$$

which imposes a constraint on the momentum uncertainty as follows

$$\frac{\hbar}{\alpha} \left[r_0 - \sqrt{r_0^2 - \alpha} \right] \leq \Delta p \leq \frac{\hbar}{\alpha} \left[r_0 + \sqrt{r_0^2 - \alpha} \right]. \quad (10)$$

So the product of Δx and Δp yields

$$\begin{aligned} \Delta x \Delta p &\geq \hbar \left[1 + \frac{\alpha}{\hbar^2} (\Delta p)^2 \right] \\ &\geq \frac{2\hbar}{\alpha} \left(r_0^2 - r_0 \sqrt{r_0^2 - \alpha} \right) = \hbar', \end{aligned} \quad (11)$$

where the second inequality is obtained by taking the lower bound of Δp . The above inequality can be rewritten as a Heisenberg-type uncertainty principle, $\Delta x \Delta p \geq \hbar'$, where $\hbar'$ may be regarded as an effective Planck constant. Thus the increase in area satisfies

$$\Delta A \geq \gamma_1 \hbar' = \frac{2\gamma_1 \hbar}{\alpha} \left(r_0^2 - r_0 \sqrt{r_0^2 - \alpha} \right). \quad (12)$$

When the particle vanishes, the information of one bit is lost and the black hole acquires the increase in entropy $(\Delta S)_{min} = \ln 2$. On the other hand, the minimum increase in the horizon area is given by the lower bound of (12), which is denoted by $(\Delta A)_{min}$. We obtain

$$\frac{dA}{dS} \simeq \frac{(\Delta A)_{min}}{(\Delta S)_{min}} = \frac{2\gamma_1 \hbar}{\alpha \ln 2} (r_0^2 - r_0 \sqrt{r_0^2 - \alpha}). \quad (13)$$

The black hole temperature (5) is deduced to

$$T \simeq \frac{\kappa}{8\pi} \cdot \frac{2\gamma_1 \hbar}{\alpha \ln 2} (r_0^2 - r_0 \sqrt{r_0^2 - \alpha}),$$

²For example, see Ref.[10]. In a rotating case, the characteristic size is represented by the black hole's irreducible mass. This point will be discussed in the next subsection.

which is not only proportional to the surface gravity but also depends on the black hole size. It should reproduce the standard result $T = \kappa/2\pi$, as $\alpha \rightarrow 0$. This requires that the calibration factor yield $\gamma_1 = 4 \ln 2$. Thus we obtain

$$T \simeq \frac{\hbar' \kappa}{2\pi}, \quad (14)$$

which is the expression for the temperature of a static and spherically symmetric black hole. Comparing the standard formula (2) with the revised version (14), we find that the latter can be obtained from the former by substituting $\hbar'$ for the Planck constant. It suggests that $\hbar'$ play the role of an effective Planck constant.

The expression (14) can be understood by reexamining the efficiency of a Geroch process. This gedanken experiment imagines a machine operating between a black hole and a remote reservoir.³ In this process, a box is filled with black body radiation from the reservoir and lowered down to the black hole surface. After emitting the radiation into the black hole, the box is moved away from the black hole. The over-all process converts heat into work with the efficiency[26]

$$\eta = 1 - \gamma_2 \kappa \ell, \quad (15)$$

where ℓ is the size of the box, and γ_2 is a coefficient factor to be determined. The smaller ℓ is, the greater η is. In practical situations, it is reasonable that the box's size is required to yield $\ell \leq 2r_0$. This is also necessary to emit the total radiation into the black hole, otherwise the photons with lower energy will not contribute to the Geroch process. On the other hand, the box must have a nonzero size, and ℓ has a minimum value which is related to the temperature of radiation, T_R . To find the relation between the temperature and efficiency, we rewrite (15) as

$$\eta = 1 - \gamma_2 (\ell T_R) \frac{\kappa}{T_R}. \quad (16)$$

For a given reservoir, the maximum value of the efficiency is determined by the smallest value for the production of ℓ and T_R . As the characteristic energy of thermal photons, the radiation temperature yields $T_R > \epsilon$, where ϵ is the photon's minimum energy which is given by the lower bound of (10). Thus we obtain

$$\begin{aligned} \ell T_R > \epsilon \ell &= \frac{\hbar}{2\alpha} \left(\ell^2 - \ell \sqrt{\ell^2 - 4\alpha} \right) \\ &\geq \frac{2\hbar}{\alpha} \left(r_0^2 - r_0 \sqrt{r_0^2 - \alpha} \right) \\ &= \hbar', \end{aligned}$$

³For details, see Ref.[26].

where we have considered $\ell \leq 2r_0$. Thus the efficiency (16) yields

$$\eta < 1 - \gamma_2 \frac{\hbar' \kappa}{T_R}. \quad (17)$$

Comparing it with the efficiency of a heat engine operating between two reservoirs, we find that the expression $\hbar' \kappa$ plays the role of the black hole temperature. This agrees with (14), up to a constant factor.

The black hole entropy can be expressed as

$$S = \int \frac{dS}{dA} dA \simeq \int \frac{(\Delta S)_{min}}{(\Delta A)_{min}} dA.$$

Considering (13) and setting $\gamma_1 = 4 \ln 2$, we obtain

$$\begin{aligned} S &\simeq \frac{1}{4} \int \frac{dA}{\hbar'} \\ &= \frac{\pi}{\hbar} \int \left(r_0 + \sqrt{r_0^2 - \alpha} \right) dr_0 \\ &= \frac{\pi}{2\hbar} \left[r_0^2 + r_0 \sqrt{r_0^2 - \alpha} - \alpha \ln(r_0 + \sqrt{r_0^2 - \alpha}) \right]. \end{aligned} \quad (18)$$

When $r_0 \gg \sqrt{\alpha}$, Bekenstein-Hawking entropy and the log-type correction, as the first two leading terms in Taylor series, are presented as

$$S = (4\hbar)^{-1} (A - \alpha \pi \ln A + \dots),$$

where the log-type correction is similar to the existing results that are derived from some concrete black holes by other methods[30]-[33].

In the context of the GUP, the heat capacity is given by

$$\begin{aligned} C &= T \frac{\partial S}{\partial T} = \frac{\hbar' \kappa}{2\pi} \cdot \frac{\partial S}{\partial A} \cdot \frac{\partial A}{\partial T} \\ &= \frac{1}{4} \left(\frac{\partial \hbar'}{\partial A} + \hbar' \kappa^{-1} \frac{\partial \kappa}{\partial A} \right)^{-1}. \end{aligned} \quad (19)$$

Direct calculation gives

$$\frac{\partial \hbar'}{\partial A} = \frac{1}{8\pi r_0} \frac{\partial \hbar'}{\partial r_0} = -\frac{\Delta \hbar}{4f}.$$

where $\Delta \hbar = \hbar' - \hbar$, $f = f(r_0) = \pi r_0 \sqrt{r_0^2 - \alpha}$. The heat capacity (19) is deduced to

$$C = C_0 f \left[\frac{\hbar'}{\hbar} f - C_0 \Delta \hbar \right]^{-1}, \quad (20)$$

where

$$C_0 = T_H \frac{\partial S_{BH}}{\partial T_H} = (4\hbar)^{-1} \kappa \frac{\partial A}{\partial \kappa},$$

is the standard heat capacity defined by the aid of Hawking temperature (2) and Bekenstein-Hawking entropy.

Let us give a remark on the temperature expression (14). In Ref.[10], Δx and Δp are identified with the black hole's radius and temperature respectively. However, this suggestion leads to a deduction that the temperature depends only on the black hole size. So the method of Ref.[10] cannot be applied to more cases with the exception of a Schwarzschild black hole. In order to explain this weakness, let us observe a Reissner-Nordström black hole in de Sitter spacetime. Its horizon radius r_0 is determined by

$$0 = F(r_0) = 1 - \frac{2M}{r_0} + \frac{Q^2}{r_0^2} - \frac{\Lambda}{3}r_0^2, \quad (21)$$

which has a very complex solution. However, we have no need to know the concrete form of r_0 , because it is useless to our discussion. Following from (21), the mass is expressed as

$$M = \frac{1}{2} \left(r_0 + \frac{Q^2}{r_0} - \frac{\Lambda}{3}r_0^3 \right), \quad (22)$$

and the surface gravity is

$$\kappa = \frac{F'(r_0)}{2} = r_0^{-1} \left(\frac{M}{r_0} - \frac{Q^2}{r_0^2} - \frac{\Lambda}{3}r_0^2 \right), \quad (23)$$

which is identified with the black hole temperature in the semiclassical framework. However, following from Ref.[10], the GUP (1) would give the temperature as follows

$$\begin{aligned} T &\sim \frac{\hbar}{\alpha} \left[r_0 - \sqrt{r_0^2 - \alpha} \right] \\ &\simeq \hbar r_0^{-1} \left(1 + \frac{3\alpha}{4r_0^2} \right), \end{aligned}$$

which cannot reproduce the standard result (as $\alpha \rightarrow 0$). Therefore, (14) is a nontrivial extension of Ref.[10], since it is suitable for more black holes and can produce the standard expression.

In the derivation of (14), a hidden assumption is that the black holes yield the laws (3) and (6), including those in de Sitter spacetime. This is an extension of Ref.[26]. There are some evidences for this assumption. For example, following from (22) and (23), we obtain

$$dM = \frac{\kappa}{8\pi} dA + \frac{Q}{r_0} dQ, \quad (24)$$

which is the first law of a Reissner-Nordström black hole in de Sitter spacetime. When the black hole captures a neutral particle, the first law becomes

$$dM = \frac{\kappa}{8\pi} dA. \quad (25)$$

On the other hand, based on a Bekenstein-type analysis, the smallest increase in the black hole mass is given by[27]

$$\Delta M \sim b\mu\kappa, \quad (26)$$

where b and μ are the particle's size and mass respectively. Considering (25) and (26), the smallest increase in horizon area is $\Delta A \sim b\mu$, which is just (6).

2.3 Kerr-Newman black hole

In Boyer-Lindquist coordinates, a Kerr-Newman black hole of mass M , charge Q and angular momentum $J = aM$ is described by

$$ds^2 = - \left(1 - \frac{2Mr - Q^2}{\rho^2} \right) dt^2 - \frac{2a(2Mr - Q^2) \sin^2 \theta}{\rho^2} dt d\phi \\ + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + \frac{\sin^2 \theta}{\rho^2} [(r^2 + a^2)^2 - a^2 \Delta \sin^2 \theta] d\phi^2,$$

where

$$\Delta = r^2 - 2Mr + a^2 + Q^2, \\ \rho^2 = r^2 + a^2 \cos^2 \theta.$$

The location of the horizon, determined by $\Delta(r_+) = 0$, is

$$r_+ = M + \sqrt{M^2 - Q^2 - a^2}. \quad (27)$$

The horizon area A , surface gravity κ , electric potential ϕ , and angular velocity Ω are respectively given by

$$A = 4\pi(r_+^2 + a^2), \\ \kappa = \frac{r_+ - M}{r_+^2 + a^2} = \frac{4\pi(r_+ - M)}{A}, \\ \phi = \frac{r_+ Q}{r_+^2 + a^2} = \frac{4\pi r_+ Q}{A}, \\ \Omega = \frac{a}{r_+^2 + a^2} = \frac{4\pi a}{A}. \quad (28)$$

These quantities yield the following relation[26]

$$dM = \frac{\kappa}{8\pi} dA + \phi dQ + \Omega dJ. \quad (29)$$

It is the first law of a Kerr-Newman black hole, in the context of mechanics.

In the previous subsection, we suggest that for a particle captured by black hole, the position uncertainty Δx yield

$$\Delta x \leq 2\rho_0, \quad (30)$$

where ρ_0 is a scale relevant to the black hole. For a static and spherically symmetric black hole, this characteristic size is identified with the twice radius of the horizon. We are confronted with a question of understanding the meaning of ρ_0 , when a rotating black hole is considered. At a first glimpse, it appears natural that ρ_0 is represented by r_+ . However, this proposal is doubtable, although it is workable for the static and spherically symmetric cases. This is because the spatial part of Boyer-Lindquist coordinates are different from ordinary polar coordinates. For instance, in a rectangular coordinates (X, Y, Z) , $r = \text{const}$ represents an ellipsoid rather than a sphere. Concretely speaking, the coordinates (r, θ, ϕ) are related to the rectangular coordinates by [34, 35, 36]

$$\begin{aligned} X &= \sqrt{r^2 + a^2} \sin \theta \cos \phi^*, \\ Y &= \sqrt{r^2 + a^2} \sin \theta \sin \phi^*, \\ Z &= r \cos \theta, \end{aligned} \quad (31)$$

where

$$\phi^* = \phi - \tan^{-1} \frac{a}{r} - a \int_{\infty}^r \frac{dr}{\Delta}.$$

Following from (31), we obtain

$$\frac{X^2 + Y^2}{r^2 + a^2} + \frac{Z^2}{r^2} = 1, \quad (32)$$

which is axially symmetric. Obviously, the surface of a Kerr-Newman black hole ($r \rightarrow r_+$) is a confocal ellipsoid. This ellipsoid is characterized by two scales: r_+ and $\sqrt{r_+^2 + a^2}$. Which is the characteristic size that represents ρ_0 ? In order to minimize ΔA , we choose the latter, i.e.

$$\rho_0 = \sqrt{r_+^2 + a^2}. \quad (33)$$

One of the evidences for (33) is that the absorption cross section for a Kerr-Newman black hole is proportional to its area [37], $\sigma_{abs} \sim A = 4\pi\rho_0^2$, which can be interpreted by the aid of a two-body process in an effective string theory that describes the collective excitations of the black hole at weak coupling [38, 39, 40]. This means that ρ_0 is indeed a characteristic size in the absorption process.

Furthermore, as argued immediately, (33) is a reasonable choice in the sense of thermodynamics, which can be explained along another line of arguments. Let us return to (30), where ρ_0 is to be determined. Replacing r_0 with ρ_0 and redoing the procedure from (9) to (13), we obtain

$$\frac{(\Delta A)_{min}}{(\Delta S)_{min}} = \frac{2\gamma_1\hbar}{\alpha \ln 2}(\rho_0^2 - \rho_0\sqrt{\rho_0^2 - \alpha}). \quad (34)$$

If ρ_0 is identified directly with r_+ , (34) becomes

$$\begin{aligned} \frac{(\Delta A)_{min}}{(\Delta S)_{min}} &= \frac{2\gamma_1\hbar}{\alpha \ln 2}(r_+^2 - r_+\sqrt{r_+^2 - \alpha}) \\ &= \frac{\gamma_1\hbar}{2\pi\alpha \ln 2} \left[A - 4\pi a^2 - \sqrt{(A - 4\pi a^2 - 2\alpha\pi)^2 - 4\alpha^2\pi^2} \right], \end{aligned}$$

which means that the entropy depends on two quantities: A and a . This contradicts Bekenstein's assumption that the entropy of a black hole is a function only of its area[26]. This would lead to a deduction incompatible with thermodynamics. Supposing $S = S(A, a)$, we have

$$\begin{aligned} dS &= \frac{\partial S}{\partial A}dA + \frac{\partial S}{\partial a}da \\ &= \frac{\partial S}{\partial A}dA + M^{-1}\frac{\partial S}{\partial a}(dJ - adM). \end{aligned} \quad (35)$$

In a reversible process, the black hole area is unchanged[41, 42], $dA = 0$. So the change in black hole mass is attributed to the work done by an external agent which changes the black hole's charge and angular momentum, and the first law (29) becomes

$$dM = \phi dQ + \Omega dJ.$$

Eq.(35) is therefore rewritten as

$$dS = M^{-1}\frac{\partial S}{\partial a}[(1 - a\Omega)dJ - a\phi dQ], \quad (36)$$

which means $dS \neq 0$ if $(\partial S/\partial a)_A \neq 0$, since Q and J are independent variables. Especially for a neutral black hole, we have

$$dS = M^{-1}\frac{r_+^2}{r_+^2 + a^2}\frac{\partial S}{\partial a}dJ.$$

The black hole entropy could be increased by an external agent which increases the angular momentum reversibly, if $(\partial S/\partial a)_A \neq 0$. This means that the entropy is not invariant in such a reversible process. This contradicts with the basic concept of thermodynamics. The crucial reason is that ρ_0 is improperly interpreted as r_+ .

What is ρ_0 ? Enlightened by the above discussion, ρ_0 should be unchanged in a reversible process. This is required by the fact that S and A are invariant in the same process. Following from (34), the black hole entropy is expressed as $S = S(A, \rho_0)$, so we have

$$dS = \frac{\partial S}{\partial A} dA + \frac{\partial S}{\partial \rho_0} d\rho_0.$$

$d\rho_0 = 0$ as $dS = dA = 0$, namely, ρ_0 is an invariant in a reversible process. For a rotating black hole in a reversible process, its irreducible mass M_{ir} is unchanged [41, 42], where

$$M_{ir} = \sqrt{\frac{A}{16\pi}} = \frac{1}{2} \sqrt{r_+^2 + a^2}. \quad (37)$$

This similarity implies that ρ_0 should be interpreted as the black hole irreducible mass, $\rho_0 \sim M_{ir}$. This can be understood in another manner. We notice that (34) involves three quantities of a black hole: the area A , entropy S and the characteristic size ρ_0 . ρ_0 is thus related not only to A but also to S . In other words, ρ_0 is a bridge which crosses the gap between A and S , hence it has geometric and thermodynamic meanings. The black hole irreducible mass agrees with this requirement. On one hand, M_{ir} is related to the area by (37). On the other hand, M_{ir} is the energy that can not be extracted by a classical process(e.g. Penrose process). In the thermodynamic sense, M_{ir} corresponds to the degraded energy that can not be transformed into work[26]. As a measure for the degradation of energy, the entropy is related to the irreducible mass by $S = S(M_{ir})$. Thus ρ_0 is endowed with geometric and thermodynamic meanings by identifying it with M_{ir} . Therefore, (33) is a natural choice in the context of thermodynamics.

Replacing r_0 with ρ_0 and doing the discussion parallel to the previous subsection, we obtain the temperature, entropy and heat capacity of a Kerr-Newman black hole, i.e.

$$\begin{aligned} T &= \frac{\hbar' \kappa}{2\pi}, \\ S &= \frac{\pi}{2\hbar} \left[\rho_0^2 + \rho_0 \sqrt{\rho_0^2 - \alpha} - \alpha \ln(\rho_0 + \sqrt{\rho_0^2 - \alpha}) \right], \\ C &= C_0 f \left[\frac{\hbar'}{\hbar} f - C_0 \Delta \hbar \right]^{-1}, \end{aligned} \quad (38)$$

where $\rho_0 = \sqrt{r_+^2 + a^2}$, and

$$\begin{aligned} \hbar' &= \frac{2\hbar}{\alpha} \left(\rho_0^2 - \rho_0 \sqrt{\rho_0^2 - \alpha} \right), \\ f &= \pi \rho_0 \sqrt{\rho_0^2 - \alpha}. \end{aligned}$$

Obviously, the expressions for these thermodynamic quantities are similar to the static and spherical black holes.

3 Black hole remnant

GUP provides a possible mechanism to prevent a black hole from complete evaporation[10]. Let us talk about our understanding of this suggestion. In the static and spherically symmetric situation, the black hole radius is restricted by GUP and must be greater than a minimum value($r_0 \geq \sqrt{\alpha}$). This restriction is necessary to (14) and (18), otherwise the two expressions would lose their physical meanings. According to (20), the black hole's heat capacity vanishes when its radius approaches the minimum value. The zero heat capacity means that no particle is emitted from the horizon, which suggests a black hole remnant. As a comparison, we temporarily ignore the GUP's impacts and first consider a Schwarzschild black hole in the standard framework. Its heat capacity is $C_0 \propto -M^2$. Due to the negative heat capacity, the black hole will evaporate to zero mass, and then $C_0 \rightarrow 0$. However, GUP makes the situation different: (i) the black hole acquires a nonzero minimum size of $\sqrt{\alpha}$; (ii) the heat capacity vanishes as $r_0 \rightarrow \sqrt{\alpha}$. In comparison with the standard result that $C_0 \rightarrow 0$ as $M \rightarrow 0$, we find that GUP elevates the black hole's "zero point energy" to a new scale determined by $r_0 = \sqrt{\alpha}$.

However, Hawking radiation makes a black hole unstable. A realistic black hole is thus non-stationary, and should be approximately described by, for example, Vaidya metric[43]

$$ds^2 = - \left[1 - \frac{2m(v)}{r} \right] dv^2 + 2dvdr + d\Omega^2,$$

where v is the Eddington-Finkelstein coordinate which denotes the retarded time. Its horizon is located by[44]-[46]

$$r_H = \frac{2m}{1 - 4\dot{m}},$$

where $\dot{m} = dm/dv$ is the mass loss rate. GUP restricts the black hole's radius by $r_H \geq \sqrt{\alpha}$, i.e.

$$\frac{2m}{1 - 4\dot{m}} \geq \sqrt{\alpha}. \quad (39)$$

For an evaporating black hole, its mass always decreases with time, i.e. $\dot{m} < 0$. Considering (39), we obtain

$$0 \leq -4\dot{m} \leq \frac{2m(v)}{\sqrt{\alpha}} - 1,$$

where $\dot{m} = 0$ denotes a black hole which stops evaporating. Obviously, $\dot{m} \rightarrow 0$ as $m(v) \rightarrow \sqrt{\alpha}/2$. Hawking radiation is shut off when the black hole evaporates to a Planck scale mass.

Black hole remnant has been suggested as an information loss reposition to resolve the black hole information problem[47, 48]. The remnant is assumed to retain the large information

of the initial black hole although it has a small size and a tiny mass. However, this idea is questionable since it violates Bekenstein's entropy bound[49], $S \leq 2\pi E\ell/\hbar$, where E denotes the energy of the system of interest and ℓ the size. Following from this bound, the remnant's information content is a few bits at most. It is too tiny to resolve the information loss problem.

Can the situation be improved when a weaker constraint is considered? In an asymptotical de Sitter spacetime, the entropy of a matter system is restricted by the so-called D bound[50]

$$S_m \leq \frac{1}{4}(A_0 - A_c), \quad (40)$$

which is derived from the generalized second law via a Geroch process, where A_0 and A_c are the areas of the cosmological horizons of pure and asymptotical de Sitter spacetimes respectively. This consideration is motivated by the astronomical observation that the current universe is dominated by the dark energy. Cosmological constant, Λ , is the simplest candidate for the dark energy. The information capacity of black hole remnant deserves to be seriously considered in the de Sitter spacetime. D-bound takes the form[50]

$$S_m \leq \pi r_g r_c, \quad (41)$$

when the gravitational radius of the matter system(r_g) is much less than the radius of the cosmological horizon(r_c). For a black hole remnant, its gravitational radius acquires the minimum value determined by GUP. Replacing $\sqrt{\alpha}$ for r_g , (41) is deduced to

$$S_r \leq \sqrt{\alpha}\pi r_c < \pi\sqrt{\frac{3\alpha}{\Lambda}}, \quad (42)$$

where we have considered $r_c < r_0 = \sqrt{3/\Lambda}$. Following from quantum statistical mechanics, the entropy bound (42) means that the number of the internal states of a black hole remnant is less than $\exp(\pi\sqrt{3\alpha/\Lambda})$. In other words, the information capacity of a black hole remnant in the de Sitter spacetime is restricted by the bound (42), which is concretely determined by the cosmological constant. In Planck units, the observed value of Λ is about $\sim 10^{-120}$, and then S_r acquires the value of 10^{60} bits at most. Comparing with Bekenstein's bound, D-bound increases the remnant's information capacity dramatically. However, the situation is still not optimistic. Considering a black hole of initial mass M_0 , its entropy is $S_0 \approx 4\pi M_0^2$, which measures the total information hidden at the moment of collapse. For a solar mass black hole, its entropy is about 10^{76} bits, which is about 16 orders greater than S_r . This means that the remnant cannot retain the total information content of the initial black hole. The discrepancy becomes more serious when the larger black holes are considered. In order for the entropy bound (42) to be workable,

the black hole mass must yield

$$M_0 < \left(\frac{3\alpha}{16\Lambda} \right)^{1/4} \sim 10^{30} m_p \sim 10^{25} g,$$

which is 8 orders less than the solar mass. Obviously, this mass scale rules out the most black holes in the universe. We therefore arrive at a conclusion that black hole remnants might not serve to resolve the information paradox.

4 Summary

This research explores an alternative expression for black hole thermodynamics in the sense of GUP (1). We try to extend the argument of Ref.[10] to more general black holes. We first consider a static and spherically symmetric black hole, and work out the temperature, entropy and heat capacity of it. These quantities are expressed by (14),(18) and (20). The similar expressions are also valid to a Kerr-Newman black hole, when r_0 is replaced by ρ_0 . For example, the temperatures of both black holes, (14) and (38), can be expressed as a unified form, $T = \hbar' \kappa / 2\pi$, if r_0 and ρ_0 are written as $\sqrt{A/4\pi}$.

Our argument is based on the analysis of a gedanken experiment that a particle is absorbed by black hole. Different from the existing literature, we suggest that the black hole irreducible mass represent the characteristic size in the absorption process, which restricts the position uncertainty of the particle falling in the black hole. This suggestion follows from two evidences: (a) the absorption cross section of a black hole is proportional to its area; (b) the entropy is an invariant in a reversible process. This suggestion can be applied to a rotating black hole and is compatible with the static and spherically symmetric black hole.

GUP (1) does not depend on the concrete form of a black hole. This implies that the temperature expression (38) may be applied to more black holes, such as the black holes with dilaton. Following from the previous discussion, this generalization should be based on two assumptions: (i) the first law (3) and the increase in area (6) are still workable; (ii) there exists a characteristic size in the absorption process, which minimizes ΔA and is an invariant in a reversible process.

Following from D-bound, a remnant cannot retain the total information of the initial black hole. Is it possible for the D-bound to be corrected by GUP? Since there are some similarities (in the sense of thermodynamics) between the cosmological horizon and black hole, we guess that a log type correction, $\ln(A_c/A_0)$, might appear in the r.h.s of (40). However, this correction is too tiny to upset the conclusion from the D-bound.

Acknowledgments

The authors would like to thank the anonymous referee who stimulates an essential improvement in this work. The authors also thank Prof. Yi Ling for a part of this work. This research is supported by NSF of China(Grant Nos.10673001, 10875057), NSF of Jiangxi province(Grant No.0612038), the key project of Chinese Ministry of Education(No.208072) and Fok Ying Tung Education Foundation(No.111008). We also acknowledge the support by the Program for Innovative Research Team of Nanchang University.

References

- [1] T. Yoneya, in *Wandering in the fields*, (World Scientific, 1987).
- [2] D. J. Gross and P. F. Mende, Nucl. Phys. B303(1988) 407.
- [3] T. Yoneya, Mod. Phys. Lett. A4(1989) 1587.
- [4] M. Maggiore, Phys. Lett. B304 (1993) 65.
- [5] L. J. Garay, Int. J. Mod. Phys. A10(1995) 145, and the references therein.
- [6] A. Kempf, G. Mangano and R.B. Mann, Phys. Rev. D52(1995) 1108.
- [7] F. Scardigli, Phys. Lett. B452(1999) 39, hep-th/9904025.
- [8] D. V. Ahluwalia, Phys. Letts. A275(2000) 31.
- [9] R. J. Adler, D. I. Santiago, Mod.Phys.Lett. A14 (1999) 1371.
- [10] R. J. Adler, P. Chen and D. I. Santiago, Gen. Rel. Grav. 33(2001) 2101; arXiv: gr-qc/0106080.
- [11] P. S. Custodio and J. E. Horvath, Class. Quant. Grav.20(2003) L197; arXiv: gr-qc/0305022.
- [12] A. E. Shalyt-Margolin, Mod. Phys. Lett. A19(2004) 2037; arXiv: gr-qc/0405069.
- [13] A. J. M. Medved and E. C. Vagenas, Phys. Rev. D70(2004) 124021, arXiv:hep-th/0411022.
- [14] B. Bolen and M. Cavaglia, Gen. Rel. Grav.37(2005) 1255; arXiv:gr-qc/0411086.
- [15] G. A. Camelia, M. Arzano, Y. Ling and G. Mandanici, Class. Quant. Grav.23(2006) 2585; arXiv:gr-qc/0506110.

- [16] M. Maziashvili, Phys. Letts. B635(2006) 232; arXiv: gr-qc/0511054.
- [17] K. Nozari and A. S. Sefiedgar, Phys. Letts. B635(2006) 156; arXiv:gr-qc/0601116.
- [18] Y. Ko, S. Lee and S. Nam, arXiv:hep-th/0608016.
- [19] Y. S. Myung, Y. W. Kim and Y.J. Park, Phys. Letts. B645(2007) 393; arXiv:gr-qc/0609031.
- [20] W. Kim, Y. W. Kim and Y. J. Park, Phys. Rev. D75(2007) 127501; arXiv: gr-qc/0702018.
- [21] K. Nouicer, Class. Quant. Grav. 24(2007) 5917; arXiv:0706.2749[gr-qc].
- [22] Mu-in Park, Phys. Letts. B 659(2008) 698; arXiv:0709.2307[hep-th].
- [23] W. Kim, E. J. Son and M. Yoon, JHEP0801(2008) 035; arXiv:0801.1439[gr-qc].
- [24] X. Han, H. R. Li and Y. Ling, Phys. Letts. B666(2008) 121; arXiv: 0807.4269[gr-qc].
- [25] F. Scardigli, arXiv:0809.1832[hep-th].
- [26] J. D. Bekenstein, Phys. Rev. D7(1973) 2333.
- [27] Z. Zhao and L. Liu, Science Bulletin 28(1983) 398.
- [28] F. Scardigli, gr-qc/0607010.
- [29] J. M. Bardeen, B. Carter and S. W. Hawking, Commun. Math. Phys.31(1973) 161.
- [30] D. N. Page, New Jour. Phys. 7(2005) 203, and the references therein.
- [31] R. Banerjee and B. R. Majhi, Phys. Lett. B662(2008) 62, arXiv:0801.0200[hep-th].
- [32] R. Banerjee, B. R. Majhi and S. Samanta, Phys. Rev. D77(2008) 124035, arXiv: 0801.3583[hep-th].
- [33] R. Banerjee and B. R. Majhi, JHEP0806(2008) 095, arXiv:0805.2220[hep-th].
- [34] R. H. Boyer and R. W. Lindquist, J.Math. Phys.8 (1967) 265.
- [35] H. C. Ohanian and R. Ruffini, *Gravitation and spacetime*(Norton,1994).
- [36] C. B. Liang, *Differential Geometry and the Foundation of General Relativity*(in chinese, Beijing Normal University Press, 2001).
- [37] J. M. Maldacena and A. Strominger, Phys. Rev. D56(1997) 4975; arxiv: hep-th/9702015.

- [38] J. M. Maldacena and A. Strominger, Phys. Rev. D55(1996) 861; arxiv: hep-th/9609026.
- [39] S. Das and S. Mathur, Nucl. Phys. B 478(1996) 561, hep-th/9606185.
- [40] M. Cvetič and F. Larsen, Phys. Rev. D56 (1997) 4994, hep-th/9705192.
- [41] D. Christodoulou, Phys. Rev. Letts. 25 (1970) 1596.
- [42] D. Christodoulou and R. Ruffini, Phys. Rev. 4 (1971) 3552.
- [43] P. C. Vaidya, Proc. Indian. Acad. Sci. A233(1951) 264.
- [44] W. A. Hiscock, Phys. Rev. D16(1977) 2673.
- [45] R. Balbinot, Phys. Rev. D33(1986) 1611.
- [46] Zhao Zheng and Dai Xian-xin, Mod. Phys. Lett. A7(1992) 1771.
- [47] J. Aharonov, A. Casher and S. Nussinov, Phys. Letts. B191(1987) 51.
- [48] J. Preskill, hep-th/9209058 and the references therein.
- [49] J. D. Bekenstein, Phys. Rev. D49(1994) 1912; arXiv:gr-qc/9307035.
- [50] R. Bousso, JHEP 0104(2001) 035; arXiv: hep-th/0012052.